

## ISDOCID: &lt;WO\_\_\_\_\_9206707A1\_I\_&gt;

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU <sup>+</sup>	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
DE*	Germany	MC	Monaco	US	United States of America
DK	Denmark				

<sup>+</sup> Any designation of "SU" has effect in the Russian Federation. It is not yet known whether any such designation has effect in other States of the former Soviet Union.

- 1 -

METHODS AND COMPOSITIONS FOR THE TREATMENT  
OF CELL PROLIFERATION DISORDERS

5 The present invention relates to methods of  
treatment of cellular proliferation disorders using  
consensus human leukocyte interferon. The invention  
also relates to pharmaceutical compositions of  
consensus human leukocyte interferon that are suitable  
for the treatment of cellular proliferation disorders.

10

Background of the Invention

Interferons are a subclass of cytokines that  
exhibit both antiviral and antiproliferative activity.  
15 On the basis of biochemical and immunological  
properties, human interferons are grouped into three  
classes: interferon-alpha (leukocyte), interferon-beta  
(fibroblast) and interferon-gamma (immune). Four  
alpha interferons (grouped into subtypes A through H)  
20 having distinct amino acid sequences have been  
identified by isolating and sequencing DNA encoding  
these polypeptides. Alpha interferons have received  
considerable attention as potential therapeutic agents  
due to their antiviral and antitumor growth inhibition.

25 The purification of interferon from human  
leukocytes isolated from the buffy coat fraction of  
whole blood is described in U.S. Patent No. 4,503,035.  
Human leukocyte interferon prepared in this manner  
contains a mixture of different human leukocyte  
30 interferon amino acid sequences. The purified material  
has a specific activity of from  $0.9 \times 10^8$ - $4 \times 10^8$   
units/mg. of protein when assayed on the MDBK bovine  
cell line and from  $2 \times 10^6$ - $7.6 \times 10^8$  units/mg. of  
protein when assayed on the Ag 1732 human cell line.  
35 The cytopathic effect inhibition assay used to

- 2 -

determine interferon anti-viral activity is disclosed in U.S. Patent No. 4,241,174. The measured interferon activity was calibrated against a reference standard for human leukocyte interferon provided by the National  
5 Institutes of Health.

The construction of recombinant DNA plasmids containing sequences encoding at least part of human leukocyte interferon and the expression in *E. coli* of a polypeptide having immunological or biological activity  
10 of human leukocyte interferon is disclosed in U.S. Patent No. 4,530,901.

The construction of hybrid alpha-interferon genes containing combinations of different subtype sequences (e.g., A and D, A and B, and A and F) is  
15 disclosed in U.S. Patent Nos. 4,414,150, 4,456,748, and 4,678,751.

U.S. Patents Nos. 4,695,623 and 4,897,471 disclose novel human leukocyte interferon polypeptides having amino acid sequences which include common or  
20 predominant amino acids found at each position among naturally-occurring alpha interferon subtype polypeptides and are referred to as consensus human leukocyte interferon (IFN-con). The IFN-con amino acid sequences disclosed are designated IFN-con<sub>1</sub>, IFN-con<sub>2</sub>,  
25 and IFN-con<sub>3</sub>. The preparation of manufactured genes encoding IFN-con and the expression of said genes in *E. coli* are also disclosed.

A purification of IFN-con<sub>1</sub> produced in *E. coli* is described in Klein et al. (*J. Chromatog.*  
30 454, 205-215 (1988)). IFN-con<sub>1</sub> purified in this manner is reported to have a specific activity of 3 x 10<sup>9</sup> units/mg. protein as measured in the cytopathic effect inhibition assay using the T98G human cell line (Fish et al. *J. Interferon Res.* 9, 97-114 (1989)).  
35 Purified IFN-con<sub>1</sub> comprises three isoforms as

- 3 -

determined by isoelectric focusing which have been identified as methionyl IFN-con<sub>1</sub>, des-methionyl IFN-con<sub>1</sub> and des-methionyl IFN-con<sub>1</sub> with its N-terminus blocked by an acetyl group. (Klein et al. Arch. Biochem. Biophys. 276, 531-537 (1990)).

Alpha-interferon is currently approved in the United States and other countries for the treatment of hairy cell leukemia, venereal warts, Kaposi's Sarcoma (a cancer commonly afflicting patients suffering from Acquired Immune Deficiency Syndrome (AIDS)), and chronic non-A, non-B hepatitis. Two variants of alpha interferon have received approval for therapeutic use: Interferon alfa-2a, marketed under the trade name ROFERON-A, and Interferon alfa-2b, marketed under the trade name INTRON-A. The amino acid sequences of ROFERON-A and INTRON-A differ at a single position but otherwise are identical to the amino acid sequence of alpha-interferon subtype 2 (subtype A).

In addition to the labeled indications, alpha-interferon is being used or evaluated alone or in conjunction with chemotherapeutic agents in a variety of other cellular proliferation disorders, including chronic myelogenous leukemia, multiple myeloma, superficial bladder cancer, skin cancers (basal cell carcinoma and malignant melanoma), renal cell carcinoma, ovarian cancer, low grade lymphocytic and cutaneous T cell lymphoma, and glioma. Alpha-interferon may be effective in combination with other chemotherapy agents for the treatment of solid tumors that arise from lung, colorectal and breast cancer (see Rosenberg et al. "Principles and Applications of Biologic Therapy" in *Cancer: Principles and Practices of Oncology*, 3rd ed., Devita et al., eds. pp. 301-547 (1989), Balmer *DICP, Ann Pharmacother* 24, 761-768 (1990)).

- 4 -

Alpha-interferons are known to affect a variety of cellular functions, including DNA replication and RNA and protein synthesis, in both normal and abnormal cells. Thus, cytotoxic effects of interferon are not restricted to tumor or virus infected cells but are also manifested in normal, healthy cells as well. As a result, undesirable side effects arise during interferon therapy, particularly when high doses are required. Administration of interferon can lead to myelosuppression resulting in reduced red blood cell, white blood cell and platelet levels. Higher doses of interferon commonly give rise to flu-like symptoms (e.g., fever, fatigue, headaches and chills), gastrointestinal disorders (e.g., anorexia, nausea and diarrhea), dizziness and coughing. It would be useful to reduce or eliminate the undesirable side effects of interferon therapy without diminishing the therapeutic benefits of such therapy.

Therefore, an object of this invention is the treatment of cell proliferation disorders such as hairy cell leukemia or Kaposi's Sarcoma with IFN-con wherein the associated undesirable side effects are diminished compared to currently practiced treatment regimens or eliminated entirely. Alternatively, an object of the invention is enhanced therapeutic benefit in the treatment of cell proliferation disorders with IFN-con compared to currently practiced regimens with no corresponding increase in the frequency or severity of undesirable side effects.

30

#### Summary of the Invention

The invention encompasses methods of treatment of a cell proliferation disorder involving administering to a mammal a therapeutically effective

35

- 5 -

amount of consensus human leukocyte interferon (IFN-con). It is shown that IFN-con has higher antiproliferative activity than INTRON-A. Therefore, treatment of a cell proliferation disorder using

5 IFN-con shows enhanced efficacy and safety compared to other currently practiced interferon treatments. In particular, the administration of a therapeutically effective amount of IFN-con results in more rapid or more extensive treatment of a cellular proliferative

10 disorder compared to currently practiced methods, wherein no corresponding increase in the frequency or severity of associated undesirable side effects occurs. In addition, a therapeutically effective amount of IFN-con may be less than said amount of an interferon

15 used in currently practiced regimens. As a result, a decreased dose of IFN-con gives the same therapeutic benefit as higher doses of other interferons but with a decrease or elimination of undesirable side effects associated with currently practiced interferon therapy.

20 IFN-con is effective in treating cell proliferation disorders frequently associated with cancer. Such disorders include, but are not limited to, hairy cell leukemia and Kaposi's Sarcoma. IFN-con may be used alone or in combination with other therapeutics

25 for the treatment of cancer and other proliferative disorders. In a preferred embodiment, IFN-con is used in conjunction with a therapeutically effective amount of one or more factors that stimulate myeloid cell proliferation or differentiation, such as granulocyte

30 colony stimulating factor (G-CSF), granulocyte/macrophage colony stimulating factor (GM-CSF), interleukin-1 (IL-1), interleukin-3 (IL-3), interleukin-6 (IL-6), erythropoietin, and stem cell factor (SCF).

- 6 -

IFN-con is a nonnaturally-occurring polypeptide having antiproliferative activity. Preferably, IFN-con is a polypeptide having the amino acid sequence of IFN-con<sub>1</sub>, IFN-con<sub>2</sub>, or IFN-con<sub>3</sub>. Most preferably, IFN-con has the amino acid sequence of IFN-con<sub>1</sub>.

The invention also relates to pharmaceutical compositions comprising a therapeutically effective amount of IFN-con along with suitable diluents, adjuvants, carriers, preservatives and/or solubilizers.

#### Brief Description of the Drawings

Figures 1 through 7 show the antiproliferative activity of IFN-con<sub>1</sub> and INTRON-A, a comparative material, on Eskol, a hairy cell leukemic cell line, when interferons were added to an Eskol cell suspension at 0.1, 0.5, 1, 5, 10, 50 and 100 ngs/ml, respectively.

Figure 8 shows the first and current median MTDs achieved by Kaposi's Sarcoma patients treated with INTRON-A, IFN-Con<sub>1</sub>, or IFN-Con<sub>1</sub> and r-metGCSF.

#### Detailed Description of the Invention

As employed herein, consensus human leukocyte interferon (IFN-con) shall mean a nonnaturally-occurring polypeptide which predominantly includes those amino acid residues which are common to all naturally-occurring human leukocyte interferon subtype sequences and which includes, at one or more of those positions wherein there is no amino acid common to all subtypes, an amino acid which predominantly occurs at that position and in no event includes any amino acid residue which is not extant in that position in at least one



- 7 -

naturally-occurring subtype. IFN-con encompasses but is not limited to the amino acid sequences designated IFN-con<sub>1</sub>, IFN-con<sub>2</sub> and IFN-con<sub>3</sub> which are disclosed in commonly owned U.S. Patents 4,695,623 and 4,897,471, the  
5 entire disclosures of which are hereby incorporated by reference. DNA sequences encoding IFN-con are synthesized as described in the above-mentioned patents.

IFN-con polypeptides are the products of expression of manufactured DNA sequences transformed or  
10 transfected into bacterial hosts, especially *E. coli*. That is, IFN-con is recombinant IFN-con. IFN-con produced in *E. coli* is purified by procedures known to those skilled in the art and generally described in Klein et al., supra (1988) for IFN-con<sub>1</sub>. Purified  
15 IFN-con may comprise a mixture of isoforms, e.g., purified IFN-con<sub>1</sub> comprises a mixture of methionyl IFN-con<sub>1</sub>, des-methionyl IFN-con<sub>1</sub> and des-methionyl IFN-con<sub>1</sub> with a blocked N-terminus (Klein et al., supra (1990)). Alternatively, IFN-con may comprise a  
20 specific, isolated isoform. Isoforms of IFN-con are separated from each other by techniques such as isoelectric focusing which are known to those skilled in the art.

The subject invention provides for a method  
25 of treating a cell proliferation disorder involving administering a therapeutically effective amount of IFN-con. A preferred embodiment of the invention is a method of treatment involving administering a therapeutically effective amount of IFN-con<sub>1</sub>, IFN-con<sub>2</sub>,  
30 or IFN-con<sub>3</sub>. Most preferably, a therapeutically effective amount of IFN-con<sub>1</sub> is administered.

IFN-con is useful for treatment of a variety of cell proliferation disorders, particularly various cancers. These disorders include, but are not limited  
35 to, hairy cell leukemia, Kaposi's Sarcoma, chronic

- 8 -

myelogenous leukemia, multiple myeloma, superficial bladder cancer, skin cancer (basal cell carcinoma and malignant melanoma), renal cell carcinoma, ovarian cancer, low grade lymphocytic and cutaneous T cell lymphoma, and glioma.

IFN-con is used alone or in combination with other therapeutics for the treatment of cancer and other proliferative disorders. IFN-con is administered in conjunction with a therapeutically effective amount of one or more chemotherapy agents such as busulfan, 5-fluorouracil (5-FU), zidovudine (AZT), leucovorin, melphalan, prednisone, cyclophosphamide, dacarbazine, cisplatin, and dipyridamole. IFN-con may also be given in conjunction with cytokines such as interleukin-2 (IL-2).

A therapeutically effective amount of IFN-con may be administered in combination with a therapeutically effective amount of one or more factors that stimulate myeloid differentiation so as to overcome the effects of myelosuppression observed during interferon treatments. Such agents include, but are not limited to, G-CSF, GM-CSF, IL-1, IL-3, IL-6, erythropoietin and SCF. Stem cell factor (SCF) stimulates the proliferation of early hematopoietic progenitor cells and has been described in U.S. Serial No. 573,616, the disclosure of which is hereby incorporated by reference.

In the working examples provided, it is shown that IFN-con<sub>1</sub> is an effective antiproliferative agent against hairy cell leukemia and AIDS-associated Kaposi's Sarcoma.

The anti-proliferative activity of IFN-con<sub>1</sub> and INTRON-A assayed on Eskol cells, a hairy cell leukemic cell line, is shown in Example 1. It is shown that IFN-con<sub>1</sub> has greater anti-proliferative activity

- 9 -

than INTRON-A over a wide range of concentrations. Similar results were obtained when IFN-con<sub>1</sub> was compared to ROFERON-A. These results indicate that IFN-con<sub>1</sub> has greater therapeutic efficacy when  
5 administered at the same concentrations as INTRON-A. Alternatively, lower concentrations of IFN-con<sub>1</sub> are required to demonstrate therapeutic efficacy equivalent to that of INTRON-A.

Example 2 describes a comparative study of  
10 IFN-Con<sub>1</sub> and INTRON-A in the treatment of AIDS-associated Kaposi's Sarcoma. It was shown that patients receiving IFN-Con<sub>1</sub> achieved higher unit doses than those patients receiving INTRON-A. In addition, patients receiving both and IFN-Con<sub>1</sub> and GCSF achieved  
15 higher doses of IFN-Con<sub>1</sub> than those patients receiving IFN-Con<sub>1</sub> alone (see Figure 8). In this study, all patients received AZT as part of their treatment of HIV infection. AZT administered alone is not effective on Kaposi's Sarcoma.

20 IFN-Con<sub>1</sub> was demonstrated to be safer than INTRON-A as judged by the reduced frequency of Grade 3 toxicity when IFN-Con<sub>1</sub> was administered. Treatment with IFN-Con<sub>1</sub> showed a reduced incidence of neutropenia and liver dysfunction compared to INTRON-A treatment  
25 while treatment with IFN-Con<sub>1</sub> and r-metGCSF completely eliminated Grade 3 toxicity (see Table 2).

Also provided for are pharmaceutical compositions comprising a therapeutically effective amount of IFN-con together with pharmaceutically  
30 acceptable carriers, adjuvants, diluents, preservatives and/or solubilizers. Pharmaceutical compositions of IFN-con include diluents of various buffers (e.g., Tris-HCl, acetate, phosphate) having a range of pH and ionic strength, carriers (e.g., human  
35 serum albumin), solubilizers (e.g., tween,

- 10 -

polysorbate), and preservatives (e.g., thimerosal, benzyl alcohol). In general, components of pharmaceutical compositions can be selected from among those commonly employed with interferons and other  
5 antiproliferative agents and which are known to those skilled in the art. A pharmaceutical composition of IFN-con is supplied as an injectable solution or as a lyophilized powder which is reconstituted in an appropriate diluent prior to injection.

10 A therapeutically effective amount of IFN-con can be determined by one skilled in the art taking into account such variables as the half-life of IFN-con preparations, route of administration and the cell proliferation disorder being tested. In general, a  
15 therapeutically effective amount of IFN-con for the treatment of a cell proliferation disorder will be in the range of  $2 \times 10^6$  to  $60 \times 10^6$  units per patient administered several times per week. Doses in the lower part of the range are effective in the treatment of  
20 hairy cell leukemia while doses in the higher part of the range are suitable for the treatment of Kaposi's Sarcoma. Therapeutically effective amounts of IFN-con will preferably result in tumor remission of 20-80% depending upon the specific tumor type for a period of  
25 at least six months.

The route of administration will preferably be by injection into the blood of a mammal where the injection may be intravenous, intramuscular, subcutaneous or intralesional. The suitability of a  
30 given pharmaceutical composition for a given route of administration will be apparent to one skilled in the art.

- 11 -

The following examples are offered to more fully illustrate the invention but are not to be construed as limiting the scope thereof.

5

## EXAMPLE 1

Anti-proliferative Activity of IFN-con<sub>1</sub> and INTRON-A

The anti-proliferative activity of IFN-con<sub>1</sub> and INTRON-A was tested on the Eskol cell line, a hairy cell leukemic cell line isolated by Dr. E. Srouf at the Indiana University Medical School. Three ml cultures of Eskol cells were incubated in RPMI medium (Gibco) at 37°C in 5% CO<sub>2</sub> containing 10% fetal calf serum for 12 hours at 1 x 10<sup>5</sup> cells/ml. IFN-con<sub>1</sub> or INTRON-A (Interferon alfa 2b; Schering Corp.) was added to a final protein concentration of 0.1 to 100 ngs/ml in 100 µl of media. The protein concentration of IFN-con<sub>1</sub> was determined by a Bradford protein assay (Bradford, Anal. Biochem. 72, 248-254 (1976)) while the concentration of INTRON-A was calculated from the specific activity (2 x 10<sup>8</sup> International units/mg protein) and unit concentration supplied by the manufacturer. The number of viable cells was determined at 24 hour intervals by exclusion of trypan blue (Sigma). 100 µl of IFN-con<sub>1</sub> or INTRON-A were added to the indicated final concentration at 24 hour intervals. Viable cell counts were an average of four independent experiments with each experiment having duplicate samples. Variation in cell counts ranged from about 5% at 24 to 48 hours to about 2% at longer time points. The results shown in Figures 1-7 are ratios of viable cell counts in the presence or absence of interferon at various times expressed as percentages.

- 12 -

The viable cell count was confirmed by measuring the incorporation of  $^3\text{H}$ -thymidine into Eskol cells incubated in the presence of IFN-con<sub>1</sub> or INTRON-A. After the 120 hour incubation period, a  
5 200  $\mu\text{l}$  cell suspension was withdrawn and incubated at  $37^\circ\text{C}$  for three hours in the presence of 5  $\mu\text{Ci/ml}$   $^3\text{H}$ -thymidine (Amersham). The cells were harvested using a Cambridge cell harvester (Cambridge  
10 Technology), washed seven times with distilled water and twice with 95% ethanol and the amount of  $^3\text{H}$ -thymidine incorporated was determined by liquid scintillation counting. The observed uptake of  $^3\text{H}$ -thymidine by Eskol cells incubated for 120 hours in the presence of IFN-con<sub>1</sub> or INTRON-A was proportional  
15 to the cell viability count.

## EXAMPLE 2

20 Safety, Tolerance and Efficacy of IFN-con<sub>1</sub>  
Administered to Patients having Kaposi's Sarcoma (KS)

A randomized, open-label study to evaluate the safety and tolerance and to define the maximum tolerated dose (MTD) of IFN-con<sub>1</sub> and INTRON-A was  
25 undertaken. IFN-con<sub>1</sub> and INTRON-A were each administered in combination with zidovudine (AZT) to patients with AIDS-associated KS. In addition, the safety, tolerance and MTD of IFN-con<sub>1</sub> was determined when administered in conjunction with AZT and E. coli  
30 produced recombinant granulocyte colony stimulating factor having a methionine residue at the amino terminal end of the polypeptide (r-metGCSF). The three treatment groups in the study were:

- 13 -

1. INTRON-A and AZT
2. IFN-con<sub>1</sub> and AZT
3. IFN-con<sub>1</sub>, AZT and r-metGCSF.

- 5 At least 12 evaluable patients are included in each treatment group.

A. Product Description

- 10 IFN-con<sub>1</sub> was produced in E. coli using methods described in U.S. Patent Nos. 4,695,623 and 4,897,471. IFN-con<sub>1</sub> was purified by procedures generally described in Klein et al., supra (1988). For subcutaneous administration in the current study,
- 15 IFN-con<sub>1</sub> was supplied as a sterile protein solution in sodium phosphate buffer. If required, dilution was made into sterile saline.

Zidovudine (AZT) was purchased from Burroughs-Wellcome Co. and used as directed on the package insert.

- 20 INTRON-A was purchased from Schering Corp. as a sterile, lyophilized formulation which was resuspended in diluent as directed on the package insert.

- r-metGCSF was produced in E. coli using methods generally described in U.S. Patent No. 4,810,643, the disclosure of which is herein incorporated by reference. r-metGCSF was prepared as a sterile protein solution in 10 mM sodium acetate, 5% mannitol and 0.004% Tween 80 at pH 4.0 at a concentration of 0.3 mg/ml. If required, dilution was
- 25 4,810,643, the disclosure of which is herein incorporated by reference. r-metGCSF was prepared as a sterile protein solution in 10 mM sodium acetate, 5% mannitol and 0.004% Tween 80 at pH 4.0 at a concentration of 0.3 mg/ml. If required, dilution was
- 30 made into sterile 5% glucose in water (D<sub>5</sub>W).

B. Dosage and Schedules

- AZT. AZT was administered to all patients at
- 35 a fixed dose of 100 mg. orally every four hours while

- 14 -

the patient is awake for a total of five doses, or 500 mg, daily.

r-metGCSF. For those patients randomized to the treatment group including r-metGCSF, doses of  
5 r-metGCSF were 1 µg/kg body weight per day, administered subcutaneously as a single bolus injection. If necessary, this dosage was increased in increments of 1 µg/kg/day (not to exceed 6 µg/kg/day) or decreased in decrements of 0.5 µg/kg/day or less, as appropriate, in  
10 order to achieve the absolute neutrophil count (ANC) target range of 5,000 to 15,000/mm<sup>3</sup>.

Interferon. Patients received either IFN-con<sub>1</sub> or INTRON-A according to a dose escalation scheme. Dosage was based upon equal units of either interferon.  
15 However, because the specific activities of the two interferons are different (2 x 10<sup>8</sup> IU/mg for INTRON-A and at least 1 x 10<sup>9</sup> IU/mg for IFN-con<sub>1</sub> as determined by the antiviral cytopathic assay described in U.S. Patent No. 4,695,623), the amount of protein by weight (in mg)  
20 at any given dose will be different for INTRON-A and IFN-con<sub>1</sub>. The dose escalation scheme used is shown below in Table 1. The dose in mg of protein corresponding to each dose level in IUs is also shown in Table 1 for each interferon.

25

30

35



- 15 -

TABLE 1

Dose Escalation Schedule for INTRON-A and IFN-Con<sub>1</sub>

5	Dose Level	Dose x 10 <sup>6</sup> IU	Dose in mg Protein	
			INTRON-A	IFN-Con <sub>1</sub>
10	1	3	0.015	0.003
	2	9	0.045	0.009
	3	12	0.060	0.012
	4	15	0.075	0.015
	5	18	0.090	0.018
15	6	21	0.105	0.021
	7	24	0.120	0.024
	8	27	0.135	0.027
	9	30	0.150	0.030

20 Patients in each of the three treatment groups shown above were administered IFN-con<sub>1</sub> or INTRON-A starting at dose level 1 daily for one week before escalating to the next highest dose level. Dose escalation occurred on days 8, 15, 22, 29, 36, 43, 50 and 57. Escalation continued until each patient reached an MTD or the maximum daily dose of 30 x 10<sup>6</sup> IUs of interferon was achieved. The MTD for an individual patient was defined as the dose level below that at which dose-limiting toxicity occurs. Toxicity was

25 graded on a scale of 0 (no toxicity) to 4 (acute toxicity) using criteria established by the World Health Organization and described further in Miller et al. (Cancer 47, 210-211 (1981)). Dose-limiting toxicity was defined as a Grade 3 or Grade 4 adverse event judged to

30 be at least possibly related to interferon. Fever and chills lasting less than 24 hours, fatigue, headache, or Grade 2 or less toxicity were not used in defining the

- 16 -

MTD unless they were determined to be intolerable to the individual patient.

At the completion of the escalation phase, patients were continued on maintenance therapy

5 consisting of daily dosing at either the patient's MTD or at the maximum dose of  $30 \times 10^6$  IUs if that was achieved. Maintenance therapy was continued until disease progression or other criteria warranted removing the patient from the study.

10 During maintenance therapy, two interferon dose reductions as a result of toxicity were permitted. After two dose reductions, no further interferon dosing modifications were allowed and patients requiring further reductions were withdrawn from the program. An

15 exception to this procedure was when the dose-limiting toxicity was neutropenia ( $ANC \leq 1000/mm^3$  on two days of an approximately one week period). In this instance, the patient was allowed to remain in the study without further reduction in interferon dose, but r-metGCSF

20 therapy was initiated at  $1 \mu g/kg$  body weight per day, administered subcutaneously, to patients not receiving r-metGCSF. For patients already in the r-metGCSF treatment group, the dose of r-metGSCF administered was escalated to the next highest level (an increase of

25  $1 \mu g/kg/day$ ).

### C. Patient Selection

A total of 49 patients have been enrolled in

30 the study. An individual is enrolled in the study only after meeting all inclusion and exclusion criteria. The significant criteria for inclusion are serologically documented HIV infection, histopathologically confirmed Kaposi's Sarcoma with measurable cutaneous or oral

35 lesion(s), acceptable immune function (as measured by

- 17 -

CD4 lymphocyte levels) and under AZT treatment for less than one year.

Among the reasons for the withdrawal of a patient from the study are a second occurrence of Grade 3 or greater toxicity during the dose escalation phase, a third occurrence of dose-limiting toxicity after the individual patient's MTD has been determined and the patient is on maintenance therapy, or a progression in KS.

10

D. Determination of MTDs for IFN-con<sub>1</sub> and INTRON-A

Using the dose escalation scheme described above for weeks 1-9 of study, followed by maintenance therapy and dose reduction when appropriate, the first and current median MTDs of INTRON-A AND IFN-Con<sub>1</sub> for the three treatment groups were determined and are shown in Figure 8. Each group consists of 15 patients. Group I (INTRON-A and AZT) attained a first MTD during dose escalation of  $9 \times 10^6$  IUs and a current MTD of  $6 \times 10^6$  IUs; Group 2 (IFN-con<sub>1</sub> and AZT) attained first and a current MTDs of  $15 \times 10^6$  IUs; and Group 3 (IFN-con<sub>1</sub>, r-metGCSF and AZT) attained first and current MTDs of  $24 \times 10^6$  IUs and  $21 \times 10^6$  IUs, respectively.

25

E. Evaluation of safety of INTRON-A and IFN-Con<sub>1</sub> treatment

The safety of INTRON-A and IFN-Con<sub>1</sub> treatment was determined by the severity of adverse effects that required interferon dose reduction. The results are summarized in Table 2.

35

- 18 -

TABLE 2

Toxicities Prompting Dose Reductions  
In Three Treatment Groups

	Frequency of Occurrence (%)			
	INTRON-A	IFN-Con <sub>1</sub> *	IFN-Con <sub>1</sub> and r-metGCSF*	
5				
10				
	Grade 2 Intolerance (Flu-like syndrome)	20	70	65
15				
	Grade 3 Neutropenia	40	10	0
20				
	Grade 3 Liver function tests	30	10	0

\*Percentages for IFN-Con<sub>1</sub> and IFN-Con<sub>1</sub> and r-metGCSF treatment groups do not add up to 100% because some patients in these groups reached the maximum dose of 30 x 10<sup>6</sup> IU with no adverse effects.

Since the study was initiated, no patients have been withdrawn as a result of toxicity clearly resulting from the administration of INTRON-A or IFN-Con<sub>1</sub>.

F. Determination of efficacy of IFN-con<sub>1</sub> and  
INTRON-A treatment

Antitumor response. Antitumor responses were assessed after four months of treatment using the AIDS Clinical Trials Group (ACTG) Oncology Committee's standard response criteria (Krown et al. *J. Clin. Oncol.* 7, 1201-1207 (1989)).

- 19 -

Immune functions. CD4 lymphocyte counts are taken every month for six months during the study to evaluate patients' immune response to HIV infection.

5           In all three treatment groups, the Kaposi's Sarcoma lesion responses and CD4 lymphocyte levels were equivalent.

10

\* \* \*

15           While the present invention has been described in terms of the preferred embodiments, it is understood that variations and modifications will occur to those skilled in the art. Therefore, it is intended that the appended claims cover all such equivalent variations which come within the scope of the invention as claimed.

- 20 -

## WHAT IS CLAIMED IS:

1. A method for the treatment of a cell proliferation disorder in a mammal comprising  
5 administering a therapeutically effective amount of consensus human leukocyte interferon.
2. A method as in Claim 1 wherein the cell proliferation disorder is hairy cell leukemia.  
10
3. A method as in Claim 1 wherein the cell proliferation disorder is Kaposi's Sarcoma.
4. A method as in Claim 1 wherein the  
15 consensus human leukocyte interferon is selected from the group consisting of IFN-con<sub>1</sub>, IFN-con<sub>2</sub>, and IFN-con<sub>3</sub>.
5. A method as in Claim 4 wherein the  
20 consensus human leukocyte interferon is IFN-con<sub>1</sub>.
6. A method as in Claim 1 wherein the consensus human leukocyte interferon is the product of procaryotic expression of an exogenous DNA sequence.  
25
7. A method as in Claim 1 wherein the therapeutically effective amount is administered by intravenous, intramuscular, subcutaneous or intralesional routes.  
30
8. A method as in Claim 1 wherein the therapeutically effective amount of consensus human leukocyte interferon is from  $2 \times 10^6$  to  $60 \times 10^6$  units per patient.  
35

- 21 -

9. A method as in Claim 1 wherein the mammal is a human.

5 10. A method as in Claim 1 further comprising administering a therapeutically effective amount of a chemotherapeutic agent.

10 11. A method as in Claim 1 further comprising administering a therapeutically effective amount of G-CSF, GM-CSF, IL-1, IL-3 or SCF.

15 12. A composition comprising a therapeutically effective amount of consensus human leukocyte interferon and a pharmaceutically acceptable diluent, adjuvant, carrier, preservative and/or solubilizer.

20 13. A composition as in Claim 12 wherein the consensus human leukocyte interferon is selected from the group consisting of IFN-con<sub>1</sub>, IFN-con<sub>2</sub>, and IFN-con<sub>3</sub>.

25 14. A composition as in Claim 13 wherein the consensus human leukocyte interferon is IFN-con<sub>1</sub>.

15. A composition as in Claim 12 wherein the consensus human leukocyte interferon is the product of procaryotic expression of an exogenous DNA sequence.

30 16. A composition as in Claim 12 which is suitable for administration by intravenous, subcutaneous, intramuscular or intralesional routes.

- 22 -


17. A composition as in Claim 12 which is supplied as an injectable solution or a lyophilized powder.

5 18. A composition as in Claim 12 further comprising a therapeutically effective amount of G-CSF, GM-CSF, IL-1, IL-3 or SCF.




FIGURE 1

Antiproliferative Activity of IFN-con1 and INTRON-A at 0.1 ng/ml




2/8  
*FIGURE 2*

Antiproliferative Activity of IFN-con1 and INTRON-A at 0.5 ng/ml



3/8

*FIGURE 3***Antiproliferative Activity of IFN-con1 and INTRON-A at 1 ng/ml**

## FIGURE 4

Antiproliferative Activity of IFN-con1 and INTRON-A at 5 ng/ml






FIGURE 5

Antiproliferative Activity of IFN-con1 and INTRON-A at 10 ng/ml




## FIGURE 6

Antiproliferative Activity of IFN-con1 and INTRON-A at 50 ng/ml




## FIGURE 7

Antiproliferative Activity of IFN-con1 and INTRON-A at 100 ng/ml



8/8

Median Maximum Tolerated Doses (MTD)(in million units) for the three treatment groups



Flavie 8



# INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US91/07722

<b>I. CLASSIFICATION OF SUBJECT MATTER</b> (if several classification symbols apply, indicate all) <sup>6</sup> According to International Patent Classification (IPC) or to both National Classification and IPC IPC(5): A61K 37/66 U.S. Class: 424/85.7														
<b>II. FIELDS SEARCHED</b> <div style="text-align: center; border-top: 1px solid black; border-bottom: 1px solid black;">Minimum Documentation Searched <sup>7</sup></div> <table style="width: 100%; border-collapse: collapse;"> <tr> <th style="width: 25%; border: 1px solid black; text-align: left;">Classification System</th> <th style="width: 75%; border: 1px solid black; text-align: left;">Classification Symbols</th> </tr> <tr> <td style="border: 1px solid black; padding: 5px;">U.S. Class</td> <td style="border: 1px solid black; padding: 5px;">424/85.7</td> </tr> </table> <div style="text-align: center; border-top: 1px solid black; border-bottom: 1px solid black;">Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched <sup>8</sup></div>			Classification System	Classification Symbols	U.S. Class	424/85.7								
Classification System	Classification Symbols													
U.S. Class	424/85.7													
Dialog-File Medicine Search terms: Interferon, consensus, proliferative														
<b>III. DOCUMENTS CONSIDERED TO BE RELEVANT <sup>9</sup></b> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 10%; border: 1px solid black;">Category <sup>9</sup></th> <th style="width: 70%; border: 1px solid black;">Citation of Document, <sup>11</sup> with indication, where appropriate, of the relevant passages <sup>12</sup></th> <th style="width: 20%; border: 1px solid black;">Relevant to Claim No. <sup>13</sup></th> </tr> </thead> <tbody> <tr> <td style="text-align: center; vertical-align: top; border: 1px solid black;">Y</td> <td style="border: 1px solid black;">US, A 4,695,523 (Stabinsky) 22 September 1987, see figure 2.</td> <td style="text-align: center; vertical-align: top; border: 1px solid black;">1-18</td> </tr> <tr> <td style="text-align: center; vertical-align: top; border: 1px solid black;">Y</td> <td style="border: 1px solid black;">Blut, Volume 53, issued 1986, T. Moritz et al., The Effect of Interferons on Cellular Defferention, " pages 361-370, see entire document.</td> <td style="text-align: center; vertical-align: top; border: 1px solid black;">1-18</td> </tr> <tr> <td style="text-align: center; vertical-align: top; border: 1px solid black;">Y</td> <td style="border: 1px solid black;">Pharmac. Ther., Volume 27, issued 1985, P&gt;B&gt; Fisher, "Effects of Interferon on Differentiation of Normal and Tumor Cells," pages 143-166, see tables 1 and 2.</td> <td style="text-align: center; vertical-align: top; border: 1px solid black;">1-18</td> </tr> </tbody> </table>			Category <sup>9</sup>	Citation of Document, <sup>11</sup> with indication, where appropriate, of the relevant passages <sup>12</sup>	Relevant to Claim No. <sup>13</sup>	Y	US, A 4,695,523 (Stabinsky) 22 September 1987, see figure 2.	1-18	Y	Blut, Volume 53, issued 1986, T. Moritz et al., The Effect of Interferons on Cellular Defferention, " pages 361-370, see entire document.	1-18	Y	Pharmac. Ther., Volume 27, issued 1985, P>B> Fisher, "Effects of Interferon on Differentiation of Normal and Tumor Cells," pages 143-166, see tables 1 and 2.	1-18
Category <sup>9</sup>	Citation of Document, <sup>11</sup> with indication, where appropriate, of the relevant passages <sup>12</sup>	Relevant to Claim No. <sup>13</sup>												
Y	US, A 4,695,523 (Stabinsky) 22 September 1987, see figure 2.	1-18												
Y	Blut, Volume 53, issued 1986, T. Moritz et al., The Effect of Interferons on Cellular Defferention, " pages 361-370, see entire document.	1-18												
Y	Pharmac. Ther., Volume 27, issued 1985, P>B> Fisher, "Effects of Interferon on Differentiation of Normal and Tumor Cells," pages 143-166, see tables 1 and 2.	1-18												
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p><sup>10</sup> Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"A" document member of the same patent family</p> </div> </div>														
<b>IV. CERTIFICATION</b> <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%; border: 1px solid black; padding: 5px;">           Date of the Actual Completion of the International Search  <div style="text-align: center; border-top: 1px solid black;">15 January 1992</div> </td> <td style="width: 50%; border: 1px solid black; padding: 5px;">           Date of Mailing of this International Search Report  <div style="text-align: center; border-top: 1px solid black; font-size: 1.2em;">10 FEB 1992</div> </td> </tr> <tr> <td style="border: 1px solid black; padding: 5px;">           International Searching Authority  <div style="text-align: center; border-top: 1px solid black;">ISA/US</div> </td> <td style="border: 1px solid black; padding: 5px;">           Signature of Authorized Officer  <div style="text-align: center; border-top: 1px solid black;">              Shelly J. Guest           </div> </td> </tr> </table>			Date of the Actual Completion of the International Search <div style="text-align: center; border-top: 1px solid black;">15 January 1992</div>	Date of Mailing of this International Search Report <div style="text-align: center; border-top: 1px solid black; font-size: 1.2em;">10 FEB 1992</div>	International Searching Authority <div style="text-align: center; border-top: 1px solid black;">ISA/US</div>	Signature of Authorized Officer <div style="text-align: center; border-top: 1px solid black;">              Shelly J. Guest           </div>								
Date of the Actual Completion of the International Search <div style="text-align: center; border-top: 1px solid black;">15 January 1992</div>	Date of Mailing of this International Search Report <div style="text-align: center; border-top: 1px solid black; font-size: 1.2em;">10 FEB 1992</div>													
International Searching Authority <div style="text-align: center; border-top: 1px solid black;">ISA/US</div>	Signature of Authorized Officer <div style="text-align: center; border-top: 1px solid black;">              Shelly J. Guest           </div>													

## FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

Y	US, A, 4,879,471 (Stabinsky) 30 January 1990, see entire document.	1-18
Y	Nature, Volume 290, issued 05 March 1981, D.V. Goeddel, et al. "The Structure of eight distinct cloned human leukocyte interferon CDNAS," pages 20-26	1-18

V. ☐ OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE<sup>1</sup>

This international search report has not been established in respect of certain claims under Article 17(2) (a) for the following reasons:

1. ☐ Claim numbers \_\_\_\_\_ because they relate to subject matter <sup>12</sup> not required to be searched by this Authority, namely:

2. ☐ Claim numbers \_\_\_\_\_, because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out <sup>12</sup>, specifically:

3. ☐ Claim numbers \_\_\_\_\_, because they are dependent claims not drafted in accordance with the second and third sentences of PCT Rule 6.4(a).

VI. ☐ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING<sup>2</sup>

This International Searching Authority found multiple inventions in this international application as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application.

2. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:

3. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers:

4. ☐ As all searchable claims could be searched without effort justifying an additional fee, the International Searching Authority did not invite payment of any additional fee.

## Remarks on Protest

- ☐ The additional search fees were accompanied by applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

CORRECTED  
VERSION \*

SECOND CORRECTED  
VERSION \*\*



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>5</sup> : <b>A61K 37/66</b>		<b>A1</b>	(11) International Publication Number: <b>WO 92/06707</b>
			(43) International Publication Date: <b>30 April 1992 (30.04.92)</b>
(21) International Application Number: <b>PCT/US91/07722</b>			(74) Agents: ODRE, Steven, M. et al.; Amgen Inc., 1840 Dehavilland Drive, Thousand Oaks, CA 91320-1789 (US).
(22) International Filing Date: <b>15 October 1991 (15.10.91)</b>			
(30) Priority data:			
599,206                      17 October 1990 (17.10.90)      US			
772,922                      15 October 1991 (15.10.91)      US			
(71) Applicant: AMGEN INC. [US/US]; 1840 Dehavilland Drive, Thousand Oaks, CA 91320-1789 (US).			(81) Designated States: AT (European patent), AU, BE (European patent), CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FI, FR (European patent), GB (European patent), GR (European patent), IT (European patent), JP, KR, LU (European patent), NL (European patent), NO, SE (European patent).
(72) Inventors: BLATT, Lawrence, M. ; 389 North Brent Street, Ventura, CA 93003 (US). TAYLOR, Milton, W. ; 3712 Brown Ridge Road, Bloomington, IN 47401 (US).			
			Published <i>With international search report.</i>

(54) Title: METHODS AND COMPOSITIONS FOR THE TREATMENT OF CELL PROLIFERATION DISORDERS

(57) Abstract

Methods for the treatment of cell proliferation disorders using consensus human leukocyte interferon are disclosed. Also disclosed are pharmaceutical compositions of consensus human leukocyte interferon.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BG	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	IE	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SK	Slovak Republic
CI	Côte d'Ivoire	LI	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	MC	Monaco	TG	Togo
DE	Germany	MG	Madagascar	UA	Ukraine
DK	Denmark	ML	Mali	US	United States of America
ES	Spain	MN	Mongolia	VN	Viet Nam
FI	Finland				

**CORRECTED  
VERSION \***

**SECOND CORRECTED  
VERSION \*\***



**PCT**

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>5</sup> :</b>  <b>A61K 37/66</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 92/06707</b>  <b>(43) International Publication Date:</b> <b>30 April 1992 (30.04.92)</b>
<b>(21) International Application Number:</b> <b>PCT/US91/07722</b> <b>(22) International Filing Date:</b> <b>15 October 1991 (15.10.91)</b>  <b>(30) Priority data:</b> 599,206                      17 October 1990 (17.10.90)      US 772,422                      15 October 1991 (15.10.91)      US  <b>(71) Applicant:</b> <b>AMGEN INC. [US/US]; 1840 Dehavilland Drive, Thousand Oaks, CA 91320-1789 (US).</b>  <b>(72) Inventors:</b> <b>BLATT, Lawrence, M. ; 389 North Brent Street, Ventura, CA 93003 (US). TAYLOR, Milton, W. ; 3712 Brown Ridge Road, Bloomington, IN 47401 (US).</b>		<b>(74) Agents:</b> <b>ODRE, Steven, M. et al.; Amgen Inc., 1840 Dehavilland Drive, Thousand Oaks, CA 91320-1789 (US).</b>  <b>(81) Designated States:</b> <b>AT (European patent), AU, BE (European patent), CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FI, FR (European patent), GB (European patent), GR (European patent), IT (European patent), JP, KR, LU (European patent), NL (European patent), NO, SE (European patent).</b>  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> <b>METHODS AND COMPOSITIONS FOR THE TREATMENT OF CELL PROLIFERATION DISORDERS</b>  <b>(57) Abstract</b>  Methods for the treatment of cell proliferation disorders using consensus human leukocyte interferon are disclosed. Also disclosed are pharmaceutical compositions of consensus human leukocyte interferon.		

\* (Referred to in PCT Gazette No. 22/1992, Section II)

\*\* (Referred to in PCT Gazette No. 16/1993, Section II)

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BG	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	IE	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SK	Slovak Republic
CI	Côte d'Ivoire	LI	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	MC	Monaco	TC	Togo
DE	Germany	MG	Madagascar	UA	Ukraine
DK	Denmark	ML	Mali	US	United States of America
ES	Spain	MN	Mongolia	VN	Viet Nam
FI	Finland				



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>5</sup> :</b> <b>A61K 37/66</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 92/06707</b> <b>(43) International Publication Date:</b> <b>30 April 1992 (30.04.92)</b>
<b>(21) International Application Number:</b> <b>PCT/US91/07722</b> <b>(22) International Filing Date:</b> <b>15 October 1991 (15.10.91)</b>  <b>(30) Priority data:</b> 599.206                      17 October 1990 (17.10.90)      US 722.922                      15 October 1991 (15.10.91)      US  <b>(71) Applicant:</b> <b>AMGEN INC. [US/US]; 1840 Dehavilland Drive, Thousand Oaks, CA 91320-1789 (US).</b>  <b>(72) Inventors:</b> <b>BLATT, Lawrence, M. ; 389 North Brent Street, Ventura, CA 93003 (US). TAYLOR, Milton, W. ; 3712 Brown Ridge Road, Bloomington, IN 47401 (US).</b>		<b>(74) Agents:</b> <b>ODRE, Steven, M. et al.; Amgen Inc., 1840 Dehavilland Drive, Thousand Oaks, CA 91320-1789 (US).</b>  <b>(81) Designated States:</b> <b>AT (European patent), AU, BE (European patent), CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FI, FR (European patent), GB (European patent), GR (European patent), IT (European patent), JP, KR, LU (European patent), NL (European patent), NO, SE (European patent).</b>  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> <b>METHODS AND COMPOSITIONS FOR THE TREATMENT OF CELL PROLIFERATION DISORDERS</b>		
<b>(57) Abstract</b>		
<p>Methods for the treatment of cell proliferation disorders using consensus human leukocyte interferon are disclosed. Also disclosed are pharmaceutical compositions of consensus human leukocyte interferon.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU <sup>+</sup>	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TC	Togo
DE <sup>+</sup>	Germany	MC	Monaco	US	United States of America
DK	Denmark				

<sup>+</sup> Any designation of "SU" has effect in the Russian Federation. It is not yet known whether any such designation has effect in other States of the former Soviet Union.



- 1 -

METHODS AND COMPOSITIONS FOR THE TREATMENT  
OF CELL PROLIFERATION DISORDERS

5 The present invention relates to methods of  
treatment of cellular proliferation disorders using  
consensus human leukocyte interferon. The invention  
also relates to pharmaceutical compositions of  
consensus human leukocyte interferon that are suitable  
for the treatment of cellular proliferation disorders.

10

Background of the Invention

Interferons are a subclass of cytokines that  
exhibit both antiviral and antiproliferative activity.  
15 On the basis of biochemical and immunological  
properties, human interferons are grouped into three  
classes: interferon-alpha (leukocyte), interferon-beta  
(fibroblast) and interferon-gamma (immune). Fourteen  
alpha interferons (grouped into subtypes A through H)  
20 having distinct amino acid sequences have been  
identified by isolating and sequencing DNA encoding  
these polypeptides. Alpha interferons have received  
considerable attention as potential therapeutic agents  
due to their antiviral and antitumor growth inhibition.

25 The purification of interferon from human  
leukocytes isolated from the buffy coat fraction of  
whole blood is described in U.S. Patent No. 4,503,035.  
Human leukocyte interferon prepared in this manner  
contains a mixture of different human leukocyte  
30 interferon amino acid sequences. The purified material  
has a specific activity of from  $0.9 \times 10^8$  -  $4 \times 10^8$   
units/mg. of protein when assayed on the MDBK bovine  
cell line and from  $2 \times 10^6$  -  $7.6 \times 10^8$  units/mg. of  
protein when assayed on the Ag 1732 human cell line.  
35 The cytopathic effect inhibition assay used to

- 2 -

determine interferon anti-viral activity is disclosed in U.S. Patent No. 4,241,174. The measured interferon activity was calibrated against a reference standard for human leukocyte interferon provided by the National  
5 Institutes of Health.

The construction of recombinant DNA plasmids containing sequences encoding at least part of human leukocyte interferon and the expression in *E. coli* of a polypeptide having immunological or biological activity  
10 of human leukocyte interferon is disclosed in U.S. Patent No. 4,530,901.

The construction of hybrid alpha-interferon genes containing combinations of different subtype sequences (e.g., A and D, A and B, and A and F) is  
15 disclosed in U.S. Patent Nos. 4,414,150, 4,456,748, and 4,678,751.

U.S. Patents Nos. 4,695,623 and 4,897,471 disclose novel human leukocyte interferon polypeptides having amino acid sequences which include common or  
20 predominant amino acids found at each position among naturally-occurring alpha interferon subtype polypeptides and are referred to as consensus human leukocyte interferon (IFN-con). The IFN-con amino acid sequences disclosed are designated IFN-con<sub>1</sub>, IFN-con<sub>2</sub>,  
25 and IFN-con<sub>3</sub>. The preparation of manufactured genes encoding IFN-con and the expression of said genes in *E. coli* are also disclosed.

A purification of IFN-con<sub>1</sub> produced in *E. coli* is described in Klein et al. (*J. Chromatog.*  
30 454, 205-215 (1988)). IFN-con<sub>1</sub> purified in this manner is reported to have a specific activity of  $3 \times 10^9$  units/mg. protein as measured in the cytopathic effect inhibition assay using the T98G human cell line (Fish et al. *J. Interferon Res.* 2, 97-114 (1989)).  
35 Purified IFN-con<sub>1</sub> comprises three isoforms as

- 3 -

determined by isoelectric focusing which have been identified as methionyl IFN-con<sub>1</sub>, des-methionyl IFN-con<sub>1</sub> and des-methionyl IFN-con<sub>1</sub> with its N-terminus blocked by an acetyl group. (Klein et al. Arch. Biochem. Biophys. 276, 531-537 (1990)).

Alpha-interferon is currently approved in the United States and other countries for the treatment of hairy cell leukemia, venereal warts, Kaposi's Sarcoma (a cancer commonly afflicting patients suffering from Acquired Immune Deficiency Syndrome (AIDS)), and chronic non-A, non-B hepatitis. Two variants of alpha interferon have received approval for therapeutic use: Interferon alfa-2a, marketed under the trade name ROFERON-A, and Interferon alfa-2b, marketed under the trade name INTRON-A. The amino acid sequences of ROFERON-A and INTRON-A differ at a single position but otherwise are identical to the amino acid sequence of alpha-interferon subtype 2 (subtype A).

In addition to the labeled indications, alpha-interferon is being used or evaluated alone or in conjunction with chemotherapeutic agents in a variety of other cellular proliferation disorders, including chronic myelogenous leukemia, multiple myeloma, superficial bladder cancer, skin cancers (basal cell carcinoma and malignant melanoma), renal cell carcinoma, ovarian cancer, low grade lymphocytic and cutaneous T cell lymphoma, and glioma. Alpha-interferon may be effective in combination with other chemotherapy agents for the treatment of solid tumors that arise from lung, colorectal and breast cancer (see Rosenberg et al. "Principles and Applications of Biologic Therapy" in *Cancer: Principles and Practices of Oncology*, 3rd ed., Devita et al., eds. pp. 301-547 (1989), Balmer *DICP, Ann Pharmacother* 24, 761-768 (1990)).

- 4 -

Alpha-interferons are known to affect a variety of cellular functions, including DNA replication and RNA and protein synthesis, in both normal and abnormal cells. Thus, cytotoxic effects of interferon are not restricted to tumor or virus infected cells but are also manifested in normal, healthy cells as well. As a result, undesirable side effects arise during interferon therapy, particularly when high doses are required. Administration of interferon can lead to myelosuppression resulting in reduced red blood cell, white blood cell and platelet levels. Higher doses of interferon commonly give rise to flu-like symptoms (e.g., fever, fatigue, headaches and chills), gastrointestinal disorders (e.g., anorexia, nausea and diarrhea), dizziness and coughing. It would be useful to reduce or eliminate the undesirable side effects of interferon therapy without diminishing the therapeutic benefits of such therapy.

Therefore, an object of this invention is the treatment of cell proliferation disorders such as hairy cell leukemia or Kaposi's Sarcoma with IFN-con wherein the associated undesirable side effects are diminished compared to currently practiced treatment regimens or eliminated entirely. Alternatively, an object of the invention is enhanced therapeutic benefit in the treatment of cell proliferation disorders with IFN-con compared to currently practiced regimens with no corresponding increase in the frequency or severity of undesirable side effects.

#### Summary of the Invention

The invention encompasses methods of treatment of a cell proliferation disorder involving administering to a mammal a therapeutically effective

- 5 -

amount of consensus human leukocyte interferon (IFN-con). It is shown that IFN-con has higher antiproliferative activity than INTRON-A. Therefore, treatment of a cell proliferation disorder using

5 IFN-con shows enhanced efficacy and safety compared to other currently practiced interferon treatments. In particular, the administration of a therapeutically effective amount of IFN-con results in more rapid or more extensive treatment of a cellular proliferative

10 disorder compared to currently practiced methods, wherein no corresponding increase in the frequency or severity of associated undesirable side effects occurs. In addition, a therapeutically effective amount of IFN-con may be less than said amount of an interferon

15 used in currently practiced regimens. As a result, a decreased dose of IFN-con gives the same therapeutic benefit as higher doses of other interferons but with a decrease or elimination of undesirable side effects associated with currently practiced interferon therapy.

20 IFN-con is effective in treating cell proliferation disorders frequently associated with cancer. Such disorders include, but are not limited to, hairy cell leukemia and Kaposi's Sarcoma. IFN-con may be used alone or in combination with other therapeutics

25 for the treatment of cancer and other proliferative disorders. In a preferred embodiment, IFN-con is used in conjunction with a therapeutically effective amount of one or more factors that stimulate myeloid cell proliferation or differentiation, such as granulocyte

30 colony stimulating factor (G-CSF), granulocyte/macrophage colony stimulating factor (GM-CSF), interleukin-1 (IL-1), interleukin-3 (IL-3), interleukin-6 (IL-6), erythropoietin, and stem cell factor (SCF).

- 6 -

IFN-con is a nonnaturally-occurring polypeptide having antiproliferative activity. Preferably, IFN-con is a polypeptide having the amino acid sequence of IFN-con<sub>1</sub>, IFN-con<sub>2</sub>, or IFN-con<sub>3</sub>. Most  
5 preferably, IFN-con has the amino acid sequence of IFN-con<sub>1</sub>.

The invention also relates to pharmaceutical compositions comprising a therapeutically effective amount of IFN-con along with suitable diluents,  
10 adjuvants, carriers, preservatives and/or solubilizers.

#### Brief Description of the Drawings

Figures 1 through 7 show the antiproliferative  
15 activity of IFN-con<sub>1</sub> and INTRON-A, a comparative material, on Eskol, a hairy cell leukemic cell line, when interferons were added to an Eskol cell suspension at 0.1, 0.5, 1, 5, 10, 50 and 100 ngs/ml, respectively.

20 Figure 8 shows the first and current median MTDs achieved by Kaposi's Sarcoma patients treated with INTRON-A, IFN-Con<sub>1</sub>, or IFN-Con<sub>1</sub> and r-metGCSF.

#### Detailed Description of the Invention

25 As employed herein, consensus human leukocyte interferon (IFN-con) shall mean a nonnaturally-occurring polypeptide which predominantly includes those amino acid residues which are common to all naturally-  
30 occurring human leukocyte interferon subtype sequences and which includes, at one or more of those positions wherein there is no amino acid common to all subtypes, an amino acid which predominantly occurs at that position and in no event includes any amino acid residue  
35 which is not extant in that position in at least one

- 7 -

naturally-occurring subtype. IFN-con encompasses but is not limited to the amino acid sequences designated IFN-con<sub>1</sub>, IFN-con<sub>2</sub> and IFN-con<sub>3</sub> which are disclosed in commonly owned U.S. Patents 4,695,623 and 4,897,471, the  
5 entire disclosures of which are hereby incorporated by reference. DNA sequences encoding IFN-con are synthesized as described in the above-mentioned patents.

IFN-con polypeptides are the products of expression of manufactured DNA sequences transformed or  
10 transfected into bacterial hosts, especially E. coli. That is, IFN-con is recombinant IFN-con. IFN-con produced in E. coli is purified by procedures known to those skilled in the art and generally described in Klein et al., supra (1988) for IFN-con<sub>1</sub>. Purified  
15 IFN-con may comprise a mixture of isoforms, e.g., purified IFN-con<sub>1</sub> comprises a mixture of methionyl IFN-con<sub>1</sub>, des-methionyl IFN-con<sub>1</sub> and des-methionyl IFN-con<sub>1</sub> with a blocked N-terminus (Klein et al., supra (1990)). Alternatively, IFN-con may comprise a  
20 specific, isolated isoform. Isoforms of IFN-con are separated from each other by techniques such as isoelectric focusing which are known to those skilled in the art.

The subject invention provides for a method  
25 of treating a cell proliferation disorder involving administering a therapeutically effective amount of IFN-con. A preferred embodiment of the invention is a method of treatment involving administering a therapeutically effective amount of IFN-con<sub>1</sub>, IFN-con<sub>2</sub>,  
30 or IFN-con<sub>3</sub>. Most preferably, a therapeutically effective amount of IFN-con<sub>1</sub> is administered.

IFN-con is useful for treatment of a variety of cell proliferation disorders, particularly various cancers. These disorders include, but are not limited  
35 to, hairy cell leukemia, Kaposi's Sarcoma, chronic

- 8 -

myelogenous leukemia, multiple myeloma, superficial bladder cancer, skin cancer (basal cell carcinoma and malignant melanoma), renal cell carcinoma, ovarian cancer, low grade lymphocytic and cutaneous T cell lymphoma, and glioma.

IFN-con is used alone or in combination with other therapeutics for the treatment of cancer and other proliferative disorders. IFN-con is administered in conjunction with a therapeutically effective amount of one or more chemotherapy agents such as busulfan, 5-fluorouracil (5-FU), zidovudine (AZT), leucovorin, melphalan, prednisone, cyclophosphamide, dacarbazine, cisplatin, and dipyrindamole. IFN-con may also be given in conjunction with cytokines such as interleukin-2 (IL-2).

A therapeutically effective amount of IFN-con may be administered in combination with a therapeutically effective amount of one or more factors that stimulate myeloid differentiation so as to overcome the effects of myelosuppression observed during interferon treatments. Such agents include, but are not limited to, G-CSF, GM-CSF, IL-1, IL-3, IL-6, erythropoietin and SCF. Stem cell factor (SCF) stimulates the proliferation of early hematopoietic progenitor cells and has been described in U.S. Serial No. 573,616, the disclosure of which is hereby incorporated by reference.

In the working examples provided, it is shown that IFN-con<sub>1</sub> is an effective antiproliferative agent against hairy cell leukemia and AIDS-associated Kaposi's Sarcoma.

The anti-proliferative activity of IFN-con<sub>1</sub> and INTRON-A assayed on Eskol cells, a hairy cell leukemic cell line, is shown in Example 1. It is shown that IFN-con<sub>1</sub> has greater anti-proliferative activity



- 9 -

than INTRON-A over a wide range of concentrations. Similar results were obtained when IFN-con<sub>1</sub> was compared to ROFERON-A. These results indicate that IFN-con<sub>1</sub> has greater therapeutic efficacy when  
5 administered at the same concentrations as INTRON-A. Alternatively, lower concentrations of IFN-con<sub>1</sub> are required to demonstrate therapeutic efficacy equivalent to that of INTRON-A.

Example 2 describes a comparative study of  
10 IFN-Con<sub>1</sub> and INTRON-A in the treatment of AIDS-associated Kaposi's Sarcoma. It was shown that patients receiving IFN-Con<sub>1</sub> achieved higher unit doses than those patients receiving INTRON-A. In addition, patients receiving both IFN-Con<sub>1</sub> and GCSF achieved  
15 higher doses of IFN-Con<sub>1</sub> than those patients receiving IFN-Con<sub>1</sub> alone (see Figure 8). In this study, all patients received AZT as part of their treatment of HIV infection. AZT administered alone is not effective on Kaposi's Sarcoma.

20 IFN-Con<sub>1</sub> was demonstrated to be safer than INTRON-A as judged by the reduced frequency of Grade 3 toxicity when IFN-Con<sub>1</sub> was administered. Treatment with IFN-Con<sub>1</sub> showed a reduced incidence of neutropenia and liver dysfunction compared to INTRON-A treatment  
25 while treatment with IFN-Con<sub>1</sub> and r-metGCSF completely eliminated Grade 3 toxicity (see Table 2).

Also provided for are pharmaceutical compositions comprising a therapeutically effective amount of IFN-con together with pharmaceutically  
30 acceptable carriers, adjuvants, diluents, preservatives and/or solubilizers. Pharmaceutical compositions of IFN-con include diluents of various buffers (e.g., Tris-HCl, acetate, phosphate) having a range of pH and ionic strength, carriers (e.g., human  
35 serum albumin), solubilizers (e.g., tween,

- 10 -

polysorbate), and preservatives (e.g., thimerosal, benzyl alcohol). In general, components of pharmaceutical compositions can be selected from among those commonly employed with interferons and other antiproliferative agents and which are known to those skilled in the art. A pharmaceutical composition of IFN-con is supplied as an injectable solution or as a lyophilized powder which is reconstituted in an appropriate diluent prior to injection.

10 A therapeutically effective amount of IFN-con can be determined by one skilled in the art taking into account such variables as the half-life of IFN-con preparations, route of administration and the cell proliferation disorder being tested. In general, a therapeutically effective amount of IFN-con for the treatment of a cell proliferation disorder will be in the range of  $2 \times 10^6$  to  $60 \times 10^6$  units per patient administered several times per week. Doses in the lower part of the range are effective in the treatment of hairy cell leukemia while doses in the higher part of the range are suitable for the treatment of Kaposi's Sarcoma. Therapeutically effective amounts of IFN-con will preferably result in tumor remission of 20-80% depending upon the specific tumor type for a period of at least six months.

The route of administration will preferably be by injection into the blood of a mammal where the injection may be intravenous, intramuscular, subcutaneous or intralesional. The suitability of a given pharmaceutical composition for a given route of administration will be apparent to one skilled in the art.

- 11 -

The following examples are offered to more fully illustrate the invention but are not to be construed as limiting the scope thereof.

5

## EXAMPLE 1

Anti-proliferative Activity of IFN-con<sub>1</sub> and INTRON-A

The anti-proliferative activity of IFN-con<sub>1</sub> and INTRON-A was tested on the Eskol cell line, a hairy cell leukemic cell line isolated by Dr. E. Srouf at the Indiana University Medical School. Three ml cultures of Eskol cells were incubated in RPMI medium (Gibco) at 37°C in 5% CO<sub>2</sub> containing 10% fetal calf serum for 12 hours at 1 x 10<sup>5</sup> cells/ml. IFN-con<sub>1</sub> or INTRON-A (Interferon alfa 2b; Schering Corp.) was added to a final protein concentration of 0.1 to 100 ngs/ml in 100 µl of media. The protein concentration of IFN-con<sub>1</sub> was determined by a Bradford protein assay (Bradford, *Anal. Biochem.* 72, 248-254 (1976)) while the concentration of INTRON-A was calculated from the specific activity (2 x 10<sup>8</sup> International units/mg protein) and unit concentration supplied by the manufacturer. The number of viable cells was determined at 24 hour intervals by exclusion of trypan blue (Sigma). 100 µl of IFN-con<sub>1</sub> or INTRON-A were added to the indicated final concentration at 24 hour intervals. Viable cell counts were an average of four independent experiments with each experiment having duplicate samples. Variation in cell counts ranged from about 5% at 24 to 48 hours to about 2% at longer time points. The results shown in Figures 1-7 are ratios of viable cell counts in the presence or absence of interferon at various times expressed as percentages.

- 12 -

The viable cell count was confirmed by measuring the incorporation of  $^3\text{H}$ -thymidine into Eskol cells incubated in the presence of IFN-con<sub>1</sub> or INTRON-A. After the 120 hour incubation period, a  
5 200  $\mu\text{l}$  cell suspension was withdrawn and incubated at  $37^\circ\text{C}$  for three hours in the presence of 5  $\mu\text{Ci/ml}$   $^3\text{H}$ -thymidine (Amersham). The cells were harvested using a Cambridge cell harvester (Cambridge  
Technology), washed seven times with distilled water  
10 and twice with 95% ethanol and the amount of  $^3\text{H}$ -thymidine incorporated was determined by liquid scintillation counting. The observed uptake of  $^3\text{H}$ -thymidine by Eskol cells incubated for 120 hours in the presence of IFN-con<sub>1</sub> or INTRON-A was proportional  
15 to the cell viability count.

## EXAMPLE 2

20 Safety, Tolerance and Efficacy of IFN-con<sub>1</sub>  
Administered to Patients having Kaposi's Sarcoma (KS)

A randomized, open-label study to evaluate the safety and tolerance and to define the maximum tolerated dose (MTD) of IFN-con<sub>1</sub> and INTRON-A was  
25 undertaken. IFN-con<sub>1</sub> and INTRON-A were each administered in combination with zidovudine (AZT) to patients with AIDS-associated KS. In addition, the safety, tolerance and MTD of IFN-con<sub>1</sub> was determined when administered in conjunction with AZT and E. coli  
30 produced recombinant granulocyte colony stimulating factor having a methionine residue at the amino terminal end of the polypeptide (r-metGCSF). The three treatment groups in the study were:

35

- 13 -

1. INTRON-A and AZT
2. IFN-con<sub>1</sub> and AZT
3. IFN-con<sub>1</sub>, AZT and r-metGCSF.

5 At least 12 evaluable patients are included in each treatment group.

A. Product Description

10 IFN-con<sub>1</sub> was produced in *E. coli* using methods described in U.S. Patent Nos. 4,695,623 and 4,897,471. IFN-con<sub>1</sub> was purified by procedures generally described in Klein et al., supra (1988). For subcutaneous administration in the current study,  
15 IFN-con<sub>1</sub> was supplied as a sterile protein solution in sodium phosphate buffer. If required, dilution was made into sterile saline.

Zidovudine (AZT) was purchased from Burroughs-Wellcome Co. and used as directed on the package insert.

20 INTRON-A was purchased from Schering Corp. as a sterile, lyophilized formulation which was resuspended in diluent as directed on the package insert.

r-metGCSF was produced in *E. coli* using methods generally described in U.S. Patent No. 4,810,643, the disclosure of which is herein  
25 incorporated by reference. r-metGCSF was prepared as a sterile protein solution in 10 mM sodium acetate, 5% mannitol and 0.004% Tween 80 at pH 4.0 at a concentration of 0.3 mg/ml. If required, dilution was  
30 made into sterile 5% glucose in water (D<sub>5</sub>W).

B. Dosage and Schedules

AZT. AZT was administered to all patients at  
35 a fixed dose of 100 mg. orally every four hours while

- 14 -

the patient is awake for a total of five doses, or 500 mg, daily.

r-metGCSF. For those patients randomized to the treatment group including r-metGCSF, doses of  
5 r-metGCSF were 1 µg/kg body weight per day, administered subcutaneously as a single bolus injection. If necessary, this dosage was increased in increments of 1 µg/kg/day (not to exceed 6 µg/kg/day) or decreased in decrements of 0.5 µg/kg/day or less, as appropriate, in  
10 order to achieve the absolute neutrophil count (ANC) target range of 5,000 to 15,000/mm<sup>3</sup>.

Interferon. Patients received either IFN-con<sub>1</sub> or INTRON-A according to a dose escalation scheme. Dosage was based upon equal units of either interferon.  
15 However, because the specific activities of the two interferons are different (2 x 10<sup>8</sup> IU/mg for INTRON-A and at least 1 x 10<sup>9</sup> IU/mg for IFN-con<sub>1</sub> as determined by the antiviral cytopathic assay described in U.S. Patent No. 4,695,623), the amount of protein by weight (in mg)  
20 at any given dose will be different for INTRON-A and IFN-con<sub>1</sub>. The dose escalation scheme used is shown below in Table 1. The dose in mg of protein corresponding to each dose level in IUs is also shown in Table 1 for each interferon.

25

30

35

- 15 -

TABLE 1

Dose Escalation Schedule for INTRON-A and IFN-Con<sub>1</sub>

5	Dose Level	Dose x 10 <sup>6</sup> IU	Dose in mg Protein	
			INTRON-A	IFN-Con <sub>1</sub>
10	1	3	0.015	0.003
	2	9	0.045	0.009
	3	12	0.060	0.012
	4	15	0.075	0.015
	5	18	0.090	0.018
15	6	21	0.105	0.021
	7	24	0.120	0.024
	8	27	0.135	0.027
	9	30	0.150	0.030

20 Patients in each of the three treatment groups shown above were administered IFN-con<sub>1</sub> or INTRON-A starting at dose level 1 daily for one week before escalating to the next highest dose level. Dose escalation occurred on days 8, 15, 22, 29, 36, 43, 50

25 and 57. Escalation continued until each patient reached an MTD or the maximum daily dose of 30 x 10<sup>6</sup> IUs of interferon was achieved. The MTD for an individual patient was defined as the dose level below that at which dose-limiting toxicity occurs. Toxicity was

30 graded on a scale of 0 (no toxicity) to 4 (acute toxicity) using criteria established by the World Health Organization and described further in Miller et al. (Cancer 47, 210-211 (1981)). Dose-limiting toxicity was defined as a Grade 3 or Grade 4 adverse event judged to

35 be at least possibly related to interferon. Fever and chills lasting less than 24 hours, fatigue, headache, or Grade 2 or less toxicity were not used in defining the

- 16 -

MTD unless they were determined to be intolerable to the individual patient.

At the completion of the escalation phase, patients were continued on maintenance therapy consisting of daily dosing at either the patient's MTD or at the maximum dose of  $30 \times 10^6$  IUs if that was achieved. Maintenance therapy was continued until disease progression or other criteria warranted removing the patient from the study.

During maintenance therapy, two interferon dose reductions as a result of toxicity were permitted. After two dose reductions, no further interferon dosing modifications were allowed and patients requiring further reductions were withdrawn from the program. An exception to this procedure was when the dose-limiting toxicity was neutropenia ( $ANC \leq 1000/mm^3$  on two days of an approximately one week period). In this instance, the patient was allowed to remain in the study without further reduction in interferon dose, but r-metGCSF therapy was initiated at  $1 \mu g/kg$  body weight per day, administered subcutaneously, to patients not receiving r-metGCSF. For patients already in the r-metGCSF treatment group, the dose of r-metGCSF administered was escalated to the next highest level (an increase of  $1 \mu g/kg/day$ ).

### C. Patient Selection

A total of 49 patients have been enrolled in the study. An individual is enrolled in the study only after meeting all inclusion and exclusion criteria. The significant criteria for inclusion are serologically documented HIV infection, histopathologically confirmed Kaposi's Sarcoma with measurable cutaneous or oral lesion(s), acceptable immune function (as measured by



- 17 -

CD4 lymphocyte levels) and under AZT treatment for less than one year.

Among the reasons for the withdrawal of a patient from the study are a second occurrence of Grade 3 or greater toxicity during the dose escalation phase, a third occurrence of dose-limiting toxicity after the individual patient's MTD has been determined and the patient is on maintenance therapy, or a progression in KS.

D. Determination of MTDs for IFN-con<sub>1</sub> and INTRON-A

Using the dose escalation scheme described above for weeks 1-9 of study, followed by maintenance therapy and dose reduction when appropriate, the first and current median MTDs of INTRON-A AND IFN-Con<sub>1</sub> for the three treatment groups were determined and are shown in Figure 8. Each group consists of 15 patients. Group I (INTRON-A and AZT) attained a first MTD during dose escalation of  $9 \times 10^6$  IUs and a current MTD of  $6 \times 10^6$  IUs; Group 2 (IFN-con<sub>1</sub> and AZT) attained first and a current MTDs of  $15 \times 10^6$  IUs; and Group 3 (IFN-con<sub>1</sub>, r-metGCSF and AZT) attained first and current MTDs of  $24 \times 10^6$  IUs and  $21 \times 10^6$  IUs, respectively.

E. Evaluation of safety of INTRON-A and IFN-Con<sub>1</sub> treatment

The safety of INTRON-A and IFN-Con<sub>1</sub> treatment was determined by the severity of adverse effects that required interferon dose reduction. The results are summarized in Table 2.

- 18 -

TABLE 2

Toxicities Prompting Dose Reductions  
In Three Treatment Groups

5	Frequency of Occurrence (%)			
	INTRON-A	IFN-Con <sub>1</sub> *	IFN-Con <sub>1</sub> and r-metGCSF*	
10	<hr/>			
	Grade 2 Intolerance (Flu-like syndrome)	20	70	65
15	Grade 3 Neutropenia	40	10	0
20	Grade 3 Liver function tests	30	10	0

25 \*Percentages for IFN-Con<sub>1</sub> and IFN-Con<sub>1</sub> and r-metGCSF treatment groups do not add up to 100% because some patients in these groups reached the maximum dose of 30 x 10<sup>6</sup> IU with no adverse effects.

30 Since the study was initiated, no patients have been withdrawn as a result of toxicity clearly resulting from the administration of INTRON-A or IFN-Con<sub>1</sub>.

F. Determination of efficacy of IFN-con<sub>1</sub> and INTRON-A treatment

35 Antitumor response. Antitumor responses were assessed after four months of treatment using the AIDS Clinical Trials Group (ACTG) Oncology Committee's standard response criteria (Krown et al. *J. Clin. Oncol.* 40 7, 1201-1207 (1989)).

- 19 -

Immune functions. CD4 lymphocyte counts are taken every month for six months during the study to evaluate patients' immune response to HIV infection.

5                    In all three treatment groups, the Kaposi's Sarcoma lesion responses and CD4 lymphocyte levels were equivalent.

10

\* \* \*

                  While the present invention has been described in terms of the preferred embodiments, it is  
15 understood that variations and modifications will occur to those skilled in the art. Therefore, it is intended that the appended claims cover all such equivalent variations which come within the scope of the invention as claimed.

- 20 -

## WHAT IS CLAIMED IS:

1. A method for the treatment of a cell proliferation disorder in a mammal comprising  
5 administering a therapeutically effective amount of consensus human leukocyte interferon.
2. A method as in Claim 1 wherein the cell proliferation disorder is hairy cell leukemia.  
10
3. A method as in Claim 1 wherein the cell proliferation disorder is Kaposi's Sarcoma.
4. A method as in Claim 1 wherein the  
15 consensus human leukocyte interferon is selected from the group consisting of IFN-con<sub>1</sub>, IFN-con<sub>2</sub>, and IFN-con<sub>3</sub>.
5. A method as in Claim 4 wherein the  
20 consensus human leukocyte interferon is IFN-con<sub>1</sub>.
6. A method as in Claim 1 wherein the consensus human leukocyte interferon is the product of procaryotic expression of an exogenous DNA sequence.  
25
7. A method as in Claim 1 wherein the therapeutically effective amount is administered by intravenous, intramuscular, subcutaneous or intralesional routes.  
30
8. A method as in Claim 1 wherein the therapeutically effective amount of consensus human leukocyte interferon is from  $2 \times 10^6$  to  $60 \times 10^6$  units per patient.  
35

- 21 -

9. A method as in Claim 1 wherein the mammal is a human.

10. A method as in Claim 1 further comprising administering a therapeutically effective amount of a chemotherapeutic agent.

11. A method as in Claim 1 further comprising administering a therapeutically effective amount of G-CSF, GM-CSF, IL-1, IL-3 or SCF.

12. A composition comprising a therapeutically effective amount of consensus human leukocyte interferon and a pharmaceutically acceptable diluent, adjuvant, carrier, preservative and/or solubilizer.

13. A composition as in Claim 12 wherein the consensus human leukocyte interferon is selected from the group consisting of IFN-con<sub>1</sub>, IFN-con<sub>2</sub>, and IFN-con<sub>3</sub>.

14. A composition as in Claim 13 wherein the consensus human leukocyte interferon is IFN-con<sub>1</sub>.

15. A composition as in Claim 12 wherein the consensus human leukocyte interferon is the product of procaryotic expression of an exogenous DNA sequence.

16. A composition as in Claim 12 which is suitable for administration by intravenous, subcutaneous, intramuscular or intralesional routes.

- 22 -

17. A composition as in Claim 12 which is supplied as an injectable solution or a lyophilized powder.

- 5           18. A composition as in Claim 12 further comprising a therapeutically effective amount of G-CSF, GM-CSF, IL-1, IL-3 or SCF.

1 / 8




FIG. 1

SUBSTITUTE SHEET

2 / 8




FIG. 2

SUBSTITUTE SHEET



3 / 8




FIG. 4

SUBSTITUTE SHEET

4 / 8




FIG. 3

SUBSTITUTE SHEET

5 / 8




FIG. 5

SUBSTITUTE SHEET

6 / 8




FIG. 6

**SUBSTITUTE SHEET**

7 / 8





FIG. 7

SUBSTITUTE SHEET

8 / 8



- ☐ Intron A®
- ☒ IFN-con1
- ☒ IFN-con1 + G-CSF

FIG. 8

**SUBSTITUTE SHEET**

# INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US91/07722

<b>I. CLASSIFICATION OF SUBJECT MATTER</b> (if several classification symbols apply, indicate all) <sup>6</sup> According to International Patent Classification (IPC) or to both National Classification and IPC IPC(5): A61K 37/66 U.S. Class: 424/85.7						
<b>II. FIELDS SEARCHED</b> <div style="text-align: center; border-top: 1px solid black; border-bottom: 1px solid black;">Minimum Documentation Searched <sup>7</sup></div> <table style="width: 100%; border-collapse: collapse;"> <tr> <th style="width: 30%; text-align: left; border-bottom: 1px solid black;">Classification System</th> <th style="text-align: left; border-bottom: 1px solid black;">Classification Symbols</th> </tr> <tr> <td style="border-bottom: 1px solid black;">U.S. Class</td> <td style="border-bottom: 1px solid black;">424/85.7</td> </tr> </table> <div style="text-align: center; border-top: 1px solid black; border-bottom: 1px solid black;">Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched <sup>8</sup></div>			Classification System	Classification Symbols	U.S. Class	424/85.7
Classification System	Classification Symbols					
U.S. Class	424/85.7					
Dialog-File Medicine Search terms: Interferon, concensus, proliferative						
<b>III. DOCUMENTS CONSIDERED TO BE RELEVANT <sup>9</sup></b>						
Category <sup>10</sup>	Citation of Document, <sup>11</sup> with indication, where appropriate, of the relevant passages <sup>12</sup>	Relevant to Claim No. <sup>13</sup>				
Y	US, A 4,695,523 (Stabinsky) 22 September 1987, see figure 2.	1-18				
Y	Blut, Volume 53, issued 1986, T. Moritz et al., The Effect of Interferons on Cellular Defferention," pages 261-370, see entire document.	1-18				
Y	Pharmac. Ther., Volume 27, issued 1985, P>9> Fisher, "Effects of Interferon on Differentiation of Normal and Tumor Cells," pages 143-166, see tables 1 and 2.	1-18				
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p><sup>14</sup> Special categories of cited documents: <sup>15</sup></p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"G" document member of the same patent family</p> </div> </div>						
<b>IV. CERTIFICATION</b>						
Date of the Actual Completion of the International Search <div style="text-align: center;">15 January 1992</div>		Date of Mailing of this International Search Report <div style="text-align: center; font-size: 1.2em; font-weight: bold;">10 FEB 1992</div>				
International Searching Authority <div style="text-align: center;">ISA/US</div>		Signature of Authorized Officer <div style="text-align: center;">             Shelly J. Guest         </div>				

FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

Y	US, A, 4,879,471 (Stabinsky) 20 January 1990, see entire document.	1-18
Y	Nature, Volume 290, issued 05 March 1981, D.V. Goeddel, et al. "The Structure of eight distinct cloned human leukocyte interferon CDNAS," pages 20-26	1-18

V. ☐ OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE<sup>1</sup>

This international search report has not been established in respect of certain claims under Article 17(2) (a) for the following reasons:

1. ☐ Claim numbers \_\_\_\_\_ because they relate to subject matter <sup>12</sup> not required to be searched by this Authority, namely:

2. ☐ Claim numbers \_\_\_\_\_, because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out <sup>12</sup>, specifically:

3. ☐ Claim numbers \_\_\_\_\_, because they are dependent claims not drafted in accordance with the second and third sentences of PCT Rule 6.4(a).

VI. ☐ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING<sup>2</sup>

This international Searching Authority found multiple inventions in this international application as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application.

2. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:

3. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers:

4. ☐ As all searchable claims could be searched without effort justifying an additional fee, the international Searching Authority did not invite payment of any additional fee.

Remark on Protest

☐ The additional search fees were accompanied by applicant's protest.

☐ No protest accompanied the payment of additional search fees.





**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**